

3. Lynn, D. E., Miller, S. G. & Oberlander, H., Establishment of a cell line from lepidopteran wing imaginal discs: Induction of newly synthesized proteins by 20-hydroxyecdysone. *Proc. Natl Acad. Sci. USA*, **79** (1982) 2589–93.
4. Lynn, D. E. & Oberlander, H., The establishment of cell lines from imaginal wing discs of *Spodoptera frugiperda* and *Plodia interpunctella*. *J. Insect Physiol.*, **29**, (1983) 591–6.
5. Oberlander, H. & Miller, S., Lepidopteran cell lines; Tools for research in physiology, development, and genetics. In *Advances in Cell Culture*, ed. K. Maramorosch. Academic Press, New York, 1987, vol. 5, pp. 187–207.
6. Dinan, L., Spindler-Barth, M. & Spindler, K.-D., Insect cell lines as tools for studying ecdysteroid action. *Invertebrate Reproduction and Development*, **18** (1990) 43–53.
7. Oberlander, H. & Ferkovich, S. M., Physiological and developmental capacities of insect cell lines. In *Insect Cell Biotechnology*, ed. K. Maramorosch & A. H. McIntosh. CRC Press, Boca Raton, Ann Arbor, London, Tokyo, 1994, pp. 127–40.
8. Ishaaya, I. & Casida, J. E., Dietary TH 6040 alters composition and enzyme activity of housefly larval cuticle. *Pestic. Biochem. Physiol.*, **4** (1974) 484–90.
9. Post, L. C., de Jong, B. J. & Vincent, W. R., 1-(2,6-Di-substituted benzoyl)-3-phenyl-urea insecticides: Inhibitors of chitin synthesis. *Pestic. Biochem. Physiol.*, **4** (1974) 473–83.
10. Oberlander, H. & Silhacek, D. L., New perspectives on the mode of action of benzoylphenyl urea insecticides. In *Insecticides with Novel Modes of Action: Mechanism and Application*, ed. I. Ishaaya & D. Degheele. Springer-Verlag, Heidelberg, 1998, in press.
11. Porcheron, P., Oberlander, H. & Leach, C. E., Ecdysteroid regulation of amino sugar uptake in a lepidopteran cell line derived from imaginal discs. *Arch. Insect. Biochem. Physiol.*, **7** (1988) 145–55.
12. Oberlander, H., Silhacek, D. L., Leach, E., Ishaaya, I. & Shaaya, E., *Arch. Insect. Biochem. Physiol.*, **18** (1991) 219–27.
13. Mikołajczyk, P., Oberlander, H., Silhacek, D. L., Ishaaya, I. & Shaaya, E., Chitin synthesis in *Spodoptera frugiperda* wing imaginal discs: I. Chlorfluazuron, diflubenzuron, and teflubenzuron inhibit incorporation but not uptake of [¹⁴C] *N*-acetyl-D-glucosamine. *Arch. Insect Biochem. Physiol.*, **25** (1994) 245–58.
14. Zimowska, G., Mikołajczyk, P., Silhacek, D. L. & Oberlander, H., Chitin synthesis in *Spodoptera frugiperda* wing imaginal discs. II. Selective action of chlorfluazuron on wheat germ agglutinin binding and cuticle ultrastructure. *Arch. Insect Biochem. Physiol.*, **27** (1994) 89–108.
15. Wing, G. D., Slawewski, R. A. & Carlson, G. R., RH 5849, A nonsteroidal ecdysone agonist: Effects on larval Lepidoptera. *Science (Washington)*, **241** (1988) 470–2.
16. Wing, D. D., RH 5849, A nonsteroidal ecdysone agonist: Effects on a *Drosophila* cell line. *Science (Washington)*, **241** (1988) 467–9.
17. Silhacek, D. L., Oberlander, H. & Porcheron, P., Action of RH 5849, a nonsteroidal ecdysteroid mimic, on *Plodia interpunctella* (Hübner) *in vivo* and *in vitro*. *Arch. Insect Biochem. Physiol.*, **15** (1990) 201–12.
18. Oberlander, H., Silhacek, D. L. & Porcheron, P., Nonsteroidal ecdysteroid agonists: Tools for the study of hormonal action. *Arch. Insect Biochem. Physiol.*, **28** (1995) 209–23.
19. Wing, K. D. & Aller, H. E., Ecdysteroid agonists as novel insect growth regulators. In *Pesticides and Alternatives*, ed. J. E. Casida. Elsevier Science Publishers, Amsterdam, Oxford, New York, 1990, pp. 251–7.
20. Smagghe, G. & Degheele, D., Biological activity and receptor binding of ecdysteroids and the ecdysteroid agonists RH 5849 and RH 5992 in imaginal wing discs of *Spodoptera exigua* (Lepidoptera: Noctuidae). *Eur. J. Entomol.*, **92** (1995) 333–40.
21. Oberlander, H. & Tomblin, C., Cuticle deposition in imaginal disks: Effects of juvenile hormone and fat body *in vitro*. *Science (Washington)*, **177** (1972) 441–2.
22. Silhacek, D. L. & Oberlander, H., Time-dosage studies of juvenile hormone action on the development of *Plodia interpunctella*. *J. Insect Physiol.*, **21** (1975) 153–61.

Enantioselective Catalysis for Agrochemicals: Synthetic Routes to (S)-Metolachlor, (R)-Metalaxyl and (αS,3R)-Clozylacon

Felix Spindler, Benoit Pugin, Hanspeter Buser, Hans-Peter Jalett, Ulrich Pittelkow & Hans-Ulrich Blaser*

Novartis Services AG, Catalysis & Synthesis Services, R-1055.6 CH-4002 Basel, Switzerland

Abstract: The application of enantioselective catalytic methods for the technical preparation of chiral agrochemicals is illustrated for three active ingredients of the acylanilide type. The key step for the technical synthesis of the herbicide (S)-metolachlor is the enantioselective hydrogenation of an imine intermediate using a novel iridium ferrocenyldiphosphine catalyst with an unprecedented high activity and 80% ee. (R)-metalaxyl and (αS,3R)-clozylacon were synthesized via the enantioselective hydrogenation of corresponding enamide precursors with Rh and Ru/binap catalysts with >95% and 99% enantiomeric purity, respectively. © 1998 Society of Chemical Industry

Pestic. Sci., **54**, 000–000 (1998)

Key words: Chiral acyl anilides; chiral switch; enantioselective hydrogenation; (R)-metalaxyl; (S)-metolachlor; (αS,3R)-clozylacon

1 Introduction

The biological properties of chiral agrochemicals are often strongly related to the absolute configuration.¹ As a consequence, the technical synthesis of pure or enriched enantiomers is of growing importance in the modern agrochemical industry. A promising technology for very efficient asymmetric syntheses is the application of chiral catalysts but, in the end, the best overall synthesis will be chosen for commercial production. For a technically feasible catalytic process, high enantioselectivity is not the only prerequisite; other factors such as catalyst productivity, catalyst activity, catalyst stability, availability and quality of the starting material, etc., can be even more important from a production point of view.² Catalyst productivity, expressed

* To whom correspondence should be addressed.
E-mail: hans-ulrich.blaser@sn.novartis.com

as turnover number (ton), determines the catalyst costs. The catalyst activity, given as average turnover frequency for high conversion (tof), determines the production capacity of existing equipment. As a rule of thumb, ton values for large-scale products should be in the order of $2\text{--}5 \times 10^4$ and tof values should be $>2 \times 10^3 \text{ h}^{-1}$. In the case of smaller volumes and/or more expensive products, the numbers can be somewhat lower. In this contribution, the development of industrially feasible enantioselective catalytic processes for the synthesis of chiral intermediates for (*S*)-metolachlor, (*R*)-metalaxyl and ($\alpha S,3R$)-clozylacon is summarized.

2 Background

Metolachlor ('Dual' Fig. 1) is one of the most important grass herbicides for use in maize. It is an *N*-chloroacetylated, *N*-alkoxyalkylated ortho-disubstituted aniline. The commercial product exists as a mixture of four stereoisomers, but about 95% of the herbicidal activity is due to the two (1*S*)-diastereomers.³ As a consequence, Dual Magnum® with (*S*)-metolachlor of 80% enantiomeric purity, introduced in 1997, has the same biological effect as the original commercial product with about 35% less environmental load.

Clozylacon (CGA 80000; Fig. 2, 2) is a fungicide especially well suited for soil application against oomycetes.¹ Four stereoisomers exist because of the stereo-

genic centre and the atropisomerism due to hindered rotation around the carbon–nitrogen bond. The desired fungicidal activity arises mainly from the isomer with the absolute configuration $\alpha S,3R$.

Unlike other fungicides racemic metalaxyl (Ridomil®) showed no phytotoxic effects against crop plants when applied for the control of *Phytophthora infestans* (Mont.) de Bary and *Pythium ultimum* (Trow.), and therefore separation of the individual enantiomers is not absolutely necessary.¹ However, because the (*R*)-enantiomer is responsible for all of the fungicidal activity, the application of enriched (*R*)-metalaxyl (Ridomil Gold®; Fig. 2, 3) is advantageous from an ecological point of view.

3 Approaches and results

3.1 (*S*)-Metolachlor. Two approaches for the enantioselective synthesis of (*S*)-metolachlor were investigated in some detail:

- The hydrogenation of the MEA enamide precursor (Fig. 3, 4) of metolachlor by analogy to the L-dopa process of Monsanto. However, all attempts to hydrogenate the three isomeric MEA enamides failed.
- The enantioselective hydrogenation of *N*-(2-ethyl-6-methylphenyl)-*N*-(1'-methoxymethyl)-ethylideneamine (MEA imine; Fig. 3, 5) turned out to be very difficult.⁴ The following catalyst parameters proved to be important: the metal, the type of the diphosphine, addition of halide ions and the presence of an acid; in addition, pressure and temperature also strongly influenced the catalyst activity.

With hydrogen at 80 bar and 50°C, using a catalyst generated *in situ* from $[\text{Ir}(\text{cod})\text{Cl}]_2$ and the ferrocenyldiphosphine ligand, xylyphos (Fig. 3, 6) (*R* = 3,5-xylyl, *R'* = Ph) and with a substrate-to-catalyst ratio of 1×10^6 , complete conversion was reached within 2–3 h with an initial tof exceeding $1.8 \times 10^6 \text{ h}^{-1}$. The best ees were $>80\%$. A production process was developed that has been in operation since the end of 1996.

3.2 ($\alpha S,3R$)-Clozylacon. Clozylacon was synthesized successfully via asymmetric hydrogenation of the enamide I (Fig. 3, 7) followed by chlorination to yield a mixture of the two atropisomers ($\alpha S,3R$)- and ($\alpha R,3R$)-clozylacon, which were separated by crystallization. Best overall results for the catalytic hydrogenation were obtained with a Ru-binap catalyst (Fig. 3, 8; 66% ee, 4000 turnovers at 100 bar and 50°C). Enantiomerically pure hydrogenation product was isolated in 63% chemical yield by a single crystallization of the crude reaction mixture.⁵

3.3 (*R*)-Metalaxyl. The key step for the enantioselective synthesis of (*R*)-metalaxyl was the enantioselective




Fig. 1. Stereoisomers of metolachlor.




Fig. 2. Structures of clozylacon and metalaxyl.




Fig. 3. Structures of intermediates and catalysts mentioned in text.

hydrogenation of the enamide II (Fig. 3, 9). Of 34 chiral Rh diphosphine catalysts tested, 12 produced ees >90% but most showed low activity. Optimization and scale-up experiments were done with the catalyst $[\text{Rh}(\text{nbd})_2]\text{BF}_4/(\text{R,R})\text{-Me-duphos}$. With hydrogen at 10 bar and 60°C, and with a substrate-to-catalyst ratio of 5×10^4 , 95.6% ee and a turnover frequency of $5.2 \times 10^3 \text{ h}^{-1}$ were obtained, well above the specified minimum limits. The alternative route via enantioselective hydrogenation of the corresponding imine turned out not to be feasible.

Acknowledgements

The summarized results are due to the efforts of several teams of very dedicated chemists, engineers and technicians and we would like to acknowledge the contributions of R. Hanreich, H.-D. Schneider, A. Togni, A. Wirth-Tijani, M. Fischer, R. Häusel, H. Landert, S. Maurer, M. Parak, G. Thoma and N. Vostenka. We also thank Rolf Bader, Beat Böhner, John Dingwall and Gerardo Ramos for their continuous encouragement and support.

References

- For problems with homochiral pesticides see Ramos Tombo, G. M. & Bellus, D., Chirality and Crop Protection. *Angew. Chem. Int. Edn. Engl.*, **30** (1991) 1193–215.
- Blaser, H. U., Pugin, B. & Spindler, F., Enantioselective synthesis. In *Applied Homogeneous Catalysis by Organometallic Complexes*, ed. B. Cornils & W. A. Herrmann. Verlag Chemie, Weinheim, 1996, pp. 992–1009.
- Moser, H., Ryhs, G. & Sauter, Hp, Der Einfluss von Atropisomerie und chiralem Zentrum auf die biologische Aktivität des Metolachlor. *Z. Naturforsch.*, **37b** (1982) 451–62.
- For a case history see Spindler, F., Pugin, B., Jalett, H. P., Buser, H. P., Pittelkow, U. & Blaser, H. U., A technically useful catalyst for the homogeneous hydrogenation of *N*-aryl imines: A case study. *Chem. Ind. (Dekker)*, **68** (1996) 153–66.
- For more details see Buser, H. P., Pugin, B., Spindler, F. & Sutter, M., Two enantioselective syntheses of a precursor of the biologically most-active isomer of CGA80 000 (Clozylacon). *Tetrahedron*, **47** (1991) 5709–16, and references therein.

Synthesis and Insecticidal Evaluation of Imidacloprid Analogs

Jérôme Boëlle,¹ Raphaël Schneider,¹ Philippe Gérardin,¹ Bernard Loubinoux,^{1*} Peter Maienfisch² & Alfred Rindlisbacher²

¹ LERMAB, Laboratoire de Chimie Organique et Microbiologie, Université Henri Poincaré, Nancy I. Faculté des Sciences, BP 239, 54506 Vandoeuvre-lès-Nancy Cedex, France

² Novartis Crop Protection AG, CH-4002 Basel, Switzerland

Abstract: Imidacloprid analogues containing a nitroalkylidene instead of a nitroguanidine unit have been prepared and evaluated for investigation as potential insecticides. No nitroalkylidene analogue showed significant activity against the test insects. © 1998 Society of Chemical Industry

Pestic. Sci., **54**, 000–000 (1998)

Key words: imidacloprid; insecticide; α -nitroalkene

1 Introduction

Imidacloprid (Fig. 1; 1) is a systemic neonicotinoid¹ insecticide discovered by Bayer.² It controls a broad range of commercially important pests such as soil and sucking insects and termites and is used as seed dressing, as a soil treatment and as a foliar treatment in such

* To whom correspondence should be addressed.
E-mail: Bernard.Loubinoux@lermab.u-nancy.fr